ème colloque de la Flotte

Virginie GAULLIER 1@, Frank CHANIER 1, Bruno VENDEVILLE 1, Gaël LYMER 1, Johanna LOFI 2, Françoise SAGE 3, Agnès MAILLARD 4, Isabelle THINON 5

The offshore-onshore project METYSS-METYSAR aims at getting a better understanding of the Miocene-Pliocene relationships between crustal tectonics, salt tectonics, and sedimentation along the East-Sardinian margin, Tyrrhenian Basin (Figure 1). In this key-area, the Tyrrhenian back-arc basin underwent recent rifting (9-5 Ma), pro parte coeval with the Messinian Salinity Crisis (MSC, 5.96-5.33 Ma), sea-floor spreading starting during Pliocene times, and post-rift reactivation. Thereby, the Tyrrhenian basin and the East-Sardinian margin are excellent candidates for studying the mechanisms of extreme lithospheric stretching and thinning, the role of pre-existing structural fabric during and after rifting, the reactivation of a passive margin and the associated deformation and sedimentation patterns during the MSC.

We use the MSC seismic markers and the deformation of viscous salt and its brittle overburden as proxies to better delineate the timing of rifting and post-rift reactivation, and especially to quantifying vertical and horizontal movements (Figure 5).




Figure 5. Idealized schematic cross section for the western Mediterranean basin, illustrating the geometric organization of the MSC seismic markers in the (Lofi et al., 2011a and 2011b). MSC: Messinian Salinity Crisis, MES: Margin Erosion Surface, BES: Bottom Erosion Surface, BS: Bottom Surface, TES: Top Erosion Surface, TS: Top Surface, UU: Upper Unit, MU: Mobile Unit, Lower Unit, CU: Complex Unit (Lofi et al., 2011).



Bassin Est-Sarde



Figure 1. Bathymetric map of the Mediterranean (from GeoMapApp) showing the study area (red rectangle).

The data set comprises

2400 km of new high-

resolution seismic pro-

the R/V "Téthys II"

(METYSS 1-2009 and

3-2011) and 4 field

trips in the Orosei area

(Figure 4).

files acquired onboard 41'30'

The Tyrrhenian Sea is a Neogene back-arc basin that opened by continental rifting and oceanic spreading related to the eastward migration

of the Apennine subduction system from middle Miocene to Pliocene times, linked to the progressive retreat of subducting African slab during convergence between Africa and Europe. Rifting of the southern Tyrrhenian Sea occurred during Serravallian-Tortonian and Messinian times, starting possibly as early as the Langhian, and was followed by sea-floor spreading in a few small deep basins from c. 5 Ma to present-day (Kastens et al., 1988; Mascle and Rehault, 1990; Jolivet and Faccenna, 2000; Sartori et al. 2001; 2004; Doglioni et al., 2004; Carminati et al., 2012) (Figure 2).




Figure 3. Geodynamic setting of the Tyrrhenian Sea. A: Geodynamic map reconstruction during Tortonian-Messinian times (6.5-5.3 Ma) (modified from Jolivet et al., 2006). The dotted bold black line shows the location of cross section C. B: Present-day geodynamic map showing the distribution of the oceanic domains and zones of stretched continental crust (modified from Jolivet et al., 2006; Faccenna et al., 2005). Dotted grey lines represent the depth of the Benioff zone, top of the descending Ionian slab (Sartori et al. 2004). The dotted bold black line shows the location of cross section D. C. and D: Schematic cross sections, respectively for Tortonian-Messinian and Present-Day times (modified from Sartori et al., 2001 and 2004; Spakman and Wortel, 2004; Jolivet et al., 2006).










- 1. LOG-UMR 8187, Univ. de Lille 1, Bâtiment SN5, Cité Scientifique, 59655 Villeneuve d'Ascq, France, virginie.gaullier@univ-lille1.fr
- 2. Géosciences Montpellier-UMR 5243, Bat. 22, Univ. Montpellier, 2 Place E. Bataillon, 34095 Montpellier Cedex 05, France
- 3. GéoAzur-UMR 7329, UPMC, Bât. 1, 250 av. A. Einstein, 06560 Valbonne, France 4. GET-OMP-Univ. P. Sabatier-UMR 5563, 14 av. E. Belin, 31400 Toulouse, France
- 5. BRGM-DGR/GBS, BP36009, 45060 Orléans Cedex 2, France



normal faults. Grey lines show the successive positions of trenches taken from kinematic within metamorphic core complexes (Jolivet et al., 2009). **B.** Structural sketch map. Orange arrows indicate the location of the two successive sea-floor spreading areas. The red dot corresponds to an cored oceanic crust of 4.3 My in age (Kastens et al., 1988).



Along the Eastern Sardinian Margin, Tortonian p.p. to Pliocene p.p. series are usually considered to be syn-rift sediments (Sartori et al., 2001; 2004) (Figure 3). Because the Messinian Salinity Crisis took place during this rifting time, the Eastern Sardinian Margin is thus a key area to document the relationships between

the evaporitic deposits of the MSC and the tectonic activity associated with the rifting process. Recent deformation is generally considered to be very weak or absent along the Eastern Sardinian margin. Study of late Pleistocene marine terraces onshore shows little recent uplift, c. + 0.04 mm/yr, restricted to Orosei area (Ferranti et al., 2006). Vertical movements within that area, which probably are not related to tectonic activity, have been attributed to a flexural response to the development of Plio-Quaternary volcanic bodies (Mariani et al., 2009). Present-day seismic activity appears to be scarce and only a few significant earthquakes (Mw ≥ 3.5, depth ≤ 50 km) have occurred recently in the northern part of the Eastern Sardinian margin. The Regional Centroid Moment Tensor (RCMT) focal mechanisms correspond to reverse faulting, revealing a surprising compressional stress regime in that area (Pondrelli et al., 2006; Nocquet, 2012).

A. MSC SEISMIC MARKERS




Figure 7. 1. "METYSS 3" seismic profile (location in yellow on Figure 6) illustrating the relationships between the MES and UU on the East-Sardinia Basin. 2. "METYSS 1" seismic profile (location in black on the middle insert) illustrating the relationships between the MES and Messinian Units (UU and MU) on the Cornaglia Terrace. 3. Distribution map of the MES, UU and MU. The distribution map of the Messinian Erosion Surface (MES) and of the Messinian units (Upper and Mobile Units) testify for the existence of a rifted basin already created by the time of the Messinian Salinity Crisis. It demonstrates therefore major pre-MSC rifting over the whole domain (Figure 7).

Our seismic data surprisingly showed that

there are no clues for Messinian syn-rift sedi-

ments along the East-Sardinia Basin and Cor-

naglia Terrace, hence no evidence for rifting

Figure 4. Color-shaded bathymetric map of the

study area (from CIESM-Ifremer Medimap Group,

2008) showing the data set used in this work.

Figure 6. Bathymetric and topographic map of the Tyrrhenian Basin and surroundings areas. The yellow, orange, white and red lines show the location of the "METYSS" seismic llustrated in the RESULTS section. ESB: East-Sardinia Basin.

——— HR seismics "METYSS 3

A. SALT TECTONICS

In un-confined basins, such as on Figures 9 and 10, salt tectonics is huge. There is apparently no basal slope of regional scale below salt. Moreover, no salt-related extensional structures, such as upslope normal faults, are visible along the basement highs that could attest for gravity gliding (Figure 11A). We therefore conclude that, there, the main driving process for generating salt structures (mainly diapirs) is local gravity spreading related to differential sediment loading (Figure 11B), or downbuilding (Figure 11C). Moreover, salt tectonics starts very early, during the UU deposition, suggesting an early differential sedimentary overload, even weak. It also means that diapirs are created by extensional process (Gaullier and Vendeville, 2005).





Figure 10. A: Part of seismic line MYS08b (Location on Figure 6) illustrating intense salt tectonics and various associated salt structures in an un-confined basin, on the Cornaglia Terrace. B: Color shaded multibeam bathymetric close up showing salt-related sea-floor deformation with isolated or aligned salt diapirs. The location corresponds to the white rectangle on Figure 6. Bathymetric data are modified from CIESM-IFREMR MediMap Group et al. (2008).

B. POST-RIFT REACTIVATION EVIDENCE BY SALT TECTONICS

B. TILTED BLOCKS AND FAN-SHAPED SEDIMENTS



Terrasse de Cornaglia

LATE PRE-MSC RIFTING Terrasse de Cornaglia

after Late Tortonian times (Figure 8). B. SYN-SALT TECTONICS FAN-SHAPED UU


> Because salt tectonics can create fan-shaped geometries in sediments, syn-rift deposits have to be carefully re-examined in order to decipher the effects of crustal tectonics (rifting) and thin-skinned salt tectonics (Figure 9).

Figure 9. Examples

of fan-shaped sedi-

ments due: A. To

crustal tectonics, i.e.



Our data show that there is no evidence for rifting processes after Late Tortonian times. Nevertheless, widespread deformation occurs during the Pliocene and is therefore attributed to post-rift reactivation. This post-rift tectonics is

characterized not only by normal faulting but also by contractional structures (See Chanier et al., this meeting). Some Pliocene vertical movements have been evidenced by discovering localized gravity gliding of the salt and its Late Messinian (UU) and Early Pliocene overburden. To the South of the study area, crustal-scale southward tilting triggered along-strike gravity gliding of salt and cover recorded by upslope extension and downslope shortening (Figure 12).

Upslope Midslope Downslope translation shortening extension

Figure 12. Example of late gravity gliding due to crustal vertical movement.

"Actions Marges" the French margins project CITS NSU Cited references: see the annex sheet.

